

The relative ages of LMC old clusters, and the case of NGC 1841

Ivo Saviane, Alfred Rosenberg, Giampaolo Piotto, & Antonio Aparicio
ESO Chile, IAC, UniPD, IAC

Abstract. Using archival HST/WFPC2 imaging of 7 LMC globular clusters, and following the methods outlined in our previous study, we have reached the tightest constrain so far on their age dispersion, which cannot be greater than ≈ 0.5 Gyr. We also confirm earlier results that their average age is comparable to that of the metal-poor Galactic globulars. Evidence is also provided that NGC 1841 is younger than the rest of LMC globulars.

1. Introduction

The LMC is the closest disk galaxy (at ~ 50 kpc) with a large ensemble of star clusters. At least 11 of them can be considered *bona fide* counterparts (e.g. Olszewski et al. 1996) of Galactic globular clusters (GGC). Their absolute ages are comparable to those of the Milky Way metal-poor ones (Brocato et al. 1996; Olsen et al. 1998; Johnson et al. 1999), however, their kinematics is strikingly different, since they show a definite disk-like rotation of $\approx 80 \text{ km s}^{-1}$ (Freeman et al. 1983). It is of great interest to understand the reasons of the simultaneous formation of the two systems in such dynamically different protogalaxies. A key input to this problem is the age dispersion of LMC globulars.

2. The Relative ages of LMC old clusters, and the case of NGC 1841

The CMDs are based on archival HST/WFPC2 data (GO 5897). Data reduction and calibration will be presented in a forthcoming paper. In Fig. 1 (left panel) the trend of $\delta(V-I)_{@2.5}$ vs. $[\text{Fe}/\text{H}]$ is compared to a set of theoretical isochrones (see R99 for details), and the six LMC clusters fall inside the ≈ 2 Gyr strip defined by metal-poor GGCs. We thus confirm that most LMC clusters are coeval with the Milky Way globulars, and more importantly, for the first time, we can constrain the age dispersion of the LMC flattened halo, which results to be as low as $\sigma_t \approx 0.5$ Gyr. According to Fig. 1 (left panel), NGC 1841 is ~ 2 Gyr younger than the mean Galactic/LMC sample, a fact that is confirmed by a direct comparison of its CMD to that of Hodge 11 (right panel of Fig. 1). Indeed, one can see that the young age is revealed by the HB-TO difference as well, since the NGC 1841 turnoff is brighter than that of Hodge 11.




Figure 1. Left panel: the run of our age-sensitive parameter vs. metallicity, for both Galactic globulars (open circles) and LMC ones (filled circles). The dotted lines represent the Vandenberg et al. (2000) models, in 1 Gyr steps (the bottommost line corresponding to 18 Gyr). Right panel: The CMD of Hodge 11 has been shifted in color and magnitude so that its RGB and HB overlap those of NGC 1841. The good match is expected from the small difference in metallicity (Palmieri et al. 2002). Fainter than $V = 21$ just a fiducial is shown for Hodge 11.

3. Conclusions

We have found that both the mean age and age dispersion of the LMC GCs is comparable to that of the metal-poor Galactic GCs, and clear evidence has been obtained in favor of a younger age for NGC 1841. Since it is the most metal-poor LMC cluster, it means that the interstellar medium (ISM) where it formed, has had a lower degree of chemical evolution. NGC 1841 could have been formed in a relatively isolated fragment of the proto-LMC, or it could have been part of an independent system now disrupted. Further support to the latter hypothesis could be the fact that NGC 1841 is the farthest cluster from the LMC center (~ 10 kpc) and that its radial velocity is incompatible with the rotation of the old LMC halo (Freeman et al. 1983). The fact that LMC globulars formed in a rotating protogalaxy with perfect synchronization with the GGCs, seems to require a joint formation scenario, possibly induced by the Milky Way early SF (e.g. Taniguchi et al. 1999).

References

- Brocato, E., et al. 1996, MNRAS, 282, 614
- Freeman, K.C., Illingworth, G., & Oemler, A. 1983, ApJ, 272, 488
- Johnson, J. A., et al. 1999, ApJ, 527, 199
- Olsen, K. A. G., et al. 1998, MNRAS, 300, 665
- Olszewski, E.W., Suntzeff, N.B., & Mateo, M. 1996, ARA&A, 34, 511
- Palmieri, R., et al. 2002, A&A, in press
- Rosenberg, A., Saviane, I., Piotto, G., & Aparicio, A. 1999, AJ, 118, 2306 (R99)
- Taniguchi, Y., Trentham, N., & Ikeuchi, S. 1999, ApJ, 526, 13
- VandenBerg, D. A., et al. 2000, ApJ, 532, 430